

## CD25 Monoclonal Antibody(Q22)

### Description

Product type	Primary Antibody
Code	POLY-MCA0292
Host	Mouse
Isotype	IgG
Size	20ul, 50ul, 100ul
Immunogen	Synthetic Peptide of CD25
Mol wt	30819
Species reactivity	Human
Clonality	Monoclonal
Recommended application	IHC-p, IF
Concentration	1 mg/ml
Full name	Interleukin-2 receptor subunit alpha
Synonyms	IL2RA; Interleukin-2 receptor subunit alpha; IL-2 receptor subunit alpha; IL-2-RA; IL-2R subunit alpha; IL2-RA; TAC antigen; p55; CD25

This product is for research use only, not for use in human, therapeutic or diagnostic procedure.

### Background


The interleukin 2 (IL2) receptor alpha (IL2RA) and beta (IL2RB) chains, together with the common gamma chain (IL2RG), constitute the high-affinity IL2 receptor. Homodimeric alpha chains (IL2RA) result in low-affinity receptor, while homodimeric beta (IL2RB) chains produce a medium-affinity receptor. Normally an integral-membrane protein, soluble IL2RA has been isolated and determined to result from extracellular proteolysis. Alternately-spliced IL2RA mRNAs have been isolated, but the significance of each is presently unknown. Mutations in this gene are associated with interleukin 2 receptor alpha deficiency.

### Recommended Dilution


IHC: 1:200

Not yet tested in other applications.

### Images



Immunohistochemical analysis of paraffin-embedded Human-liver tissue. 1. CD25 Monoclonal antibody(Q22) was diluted at 1:200(4°C, overnight). 2. Sodium citrate pH 6.0 was used for antibody retrieval(>98°C, 20min). 3. Secondary antibody was diluted at 1:200(room temperature, 30min). Negative control was used by secondary antibody only.



Immunohistochemical analysis of paraffin-embedded Mouse Spleen Tissue using CD 25 Mouse Monoclonal antibody diluted at 1:500.

Storage

-20°C for one year

Poly-Dtech, 204 Avenue de Colmar, Strasbourg, France

[contact@poly-dtech.com](mailto:contact@poly-dtech.com) | [www.poly-dtech.com](http://www.poly-dtech.com)